

# Journal of Child, Adult Vaccines and Immunology

Volume - 6, Issue - 1

Research Article

Published Date:-2022-09-02 12:11:26

[Follow-up of patients with sarcoidosis in an internal medicine unit from a hospital in Asturias, Spain. Analysis of extrapulmonary manifestations](#)

Sarcoidosis is a multisystem disease of unknown origin that is characterized by non-caseating epithelioid granuloma formation within various organs, mainly the lungs [1].

The objective of this study is the analysis of sarcoidosis patients affected by pulmonary manifestation alone or extrapulmonary expression.

---

Brief Communication

Published Date:-2022-09-02 11:48:58

[Impact of coronavirus pandemic on safety and time of administration of subcutaneous immunotherapy among pediatric patients](#)

Introduction: Allergen immunotherapy is the only targeted therapy that can modify the natural course of allergic diseases. In pediatric patients, SCIT with aeroallergens is an effective treatment and should be considered as a preventive strategy in the treatment of allergic diseases, even though one of the major concerns about it is its safety. The main purposes of this study were to assess the safety of SCIT ultra-rush schedules with polymerized extracts in a pediatric population and to determine the impact of the COVID-19 pandemic on the safety and time of administration of subcutaneous immunotherapy among pediatric patients.

Methods: A retrospective medical records review of patients under 18 years of age undergoing SCIT was made and re-scheduling due to restrictions imposed by the COVID-19 pandemic was recorded.

Results: A total of 192 pediatric patients were included. Fifty-nine (31%) had local reactions and systemic reactions were not reported. In March 2020, the first case of COVID-19 was diagnosed in Portugal and all non-urgent appointments and procedures were postponed. In our group of pediatric patients, 43 (22%) were referred to primary care, 38 (20%) stopped AIT definitively and 111 (58%) maintained administrations in the hospital. Only 2 (2%) of them had reactions upon reinitiation.

Conclusion: In this study, the ultra-rush protocol using polymerized extracts was safe in pediatric patients. Although the effectiveness of AIT may be compromised due to prolonged suspension of the treatment, it is important to note that despite longer interruptions, administrations may continue without compromising safety, maintaining shorter visits and a lower number of injections.

---